$$X(5568)^{\pm}$$

$$I(J^P) = ?(?^?)$$

OMITTED FROM SUMMARY TABLE

Seen as a peak in the $B_s\pi^\pm$ mass spectrum with a significance of more than 3σ by ABAZOV 16E and ABAZOV 18A in inclusive $p\overline{p}$ collisions at 1.96 TeV. Not seen by AAIJ 16AI, AABOUD 18L, AALTONEN 18A, and SIRUNYAN 18J. Needs confirmation.

$X(5568)^{\pm}$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID T		TECN	COMMENT
5566.9 ^{+3.2} +0.6	278	¹ ABAZOV	18A	D0	$p\overline{p} \rightarrow B_s^0 \pi^{\pm} X$

• • • We do not use the following data for averages, fits, limits, etc. • •

133

133

$$5567.8 \pm 2.9 ^{+0.9}_{-1.9}$$

² ABAZOV 16E D0
$$p\overline{p} \rightarrow B_s^0 \pi^{\pm} X$$

$X(5568)^{\pm}$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID		TECN	COMMENT
18.6 ^{+7.9} +3.5 -6.1-3.8	278	¹ ABAZOV	18A	D0	$p\overline{p} \rightarrow B_{S}\pi^{\pm}X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$$21.9 \pm 6.4 ^{+5.0}_{-2.5}$$

ABAZOV

16E D0 $p \overline{p} \rightarrow B_s \pi^{\pm} X$

$X(5568)^{\pm}$ DECAY MODES

	Mode	Fraction (Γ_i/Γ)
$\overline{\Gamma_1}$	$B_{s}\pi^{\pm}$	seen

$\Gamma(B_s\pi^\pm)/\Gamma_{total}$							Γ_1/Γ
VALUE	EVTS	DOCUMENT ID		TECN	COMME	NT	
seen	145	$^{ m 1}$ ABAZOV	18A	D0	$p\overline{p} o$	$B_{s}^{0}\pi^{\pm}X$ $B_{s}^{0}\pi^{\pm}X$	
seen	133	² ABAZOV	16E	D0	$p\overline{p} \rightarrow$	$B_{s}^{0}\pi^{\pm}X$	
• • We do not use the following data for averages, fits, limits, etc. • •							
not seen		³ AABOUD ⁴ AALTONEN ⁵ SIRUNYAN ⁶ AAIJ	18L	ATLS	$pp \rightarrow$	$B_s^0 \pi^{\pm} X$	
not seen		⁴ AALTONEN	18A	CDF	$p\overline{p} o$	$B_s^0 \pi^{\pm} X$	
not seen		⁵ SIRUNYAN	18J	CMS	pp o	$B_s^0 \pi^{\pm} X$	
not seen		⁶ AAIJ	16AI	LHCB	pp o	$B_s^0 \pi^{\pm} X$	
						3	

https://pdg.lbl.gov

Page 1

Created: 6/1/2021 08:32

¹ From the combined analysis of $B_s^0 \to J/\psi \phi$ and $B_s^0 \to D_s^\pm \mu^\mp X$ decays. ² Assumes $X(5568)^\pm \to B_s \pi^\pm$ decay. If $X(5568)^\pm \to B_s^* \pi^\pm$ decay is assumed, the mass shifts upward by 49 MeV.

¹ From the combined analysis of $B^0_s \to J/\psi \phi$ and $B^0_s \to D^\pm_s \mu^\mp X$ decays.

$X(5568)^{\pm}$ REFERENCES

AABOUD	18L	PRL 120 202007	M. Aaboud et al.	(ATLAS Collab.)
AALTONEN	18A	PRL 120 202006	T. Aaltonen et al.	(CDF Collab.)
ABAZOV	18A	PR D97 092004	V.M. Abazov <i>et al.</i>	(D0 Collab.)
SIRUNYAN	18J	PRL 120 202005	A.M. Sirunyan et al.	(CMS Collab.)
AAIJ	16AI	PRL 117 152003	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABAZOV	16E	PRL 117 022003	V.M. Abazov et al.	(D0 Collab.)

Created: 6/1/2021 08:32

¹With B_s mesons reconstructed in decays to $D_s^{\pm} \mu^{\mp} X$.

² Seen in $p\overline{p}$ collisions at 1.96 TeV at a rate of $(8.6\pm1.9\pm1.4)\%$ relative to inclusive B_{S} production in the kinematic region $10 < p_{T}(B_{S}) < 30$ GeV/c, with B_{S} mesons reconstructed in decays to $J/\psi \phi$. An alternative possibility, $X(5568)^{\pm} \rightarrow B_{S}^{*} \pi^{\pm}$ with a missing γ , could not be ruled out.

 $^{^3}$ Not seen in 24.4 fb $^{-1}$ of pp collision data at $\sqrt{s}=7$ and 8 TeV with $B_{\rm S}$ mesons reconstructed in decays to $J/\psi\phi$. An upper limit on the production rate times branching fraction for $X(5568)^\pm\to B_{\rm S}\,\pi^\pm$ relative to inclusive $B_{\rm S}$ production is less than 1.5% at $p_T(B_{\rm S})>10$ GeV/c and less than 1.6% at $p_T(B_{\rm S})>15$ GeV/c at 95% CL.

⁴ Not seen in 9.6 fb⁻¹ of $p\overline{p}$ collision data at $\sqrt{s}=1.96$ TeV with $B_{\mathcal{S}}$ mesons reconstructed in decays to $J/\psi\,\phi$. An upper limit on the production rate times branching fraction for $X(5568)^\pm\to~B_{\mathcal{S}}\,\pi^\pm$ relative to inclusive $B_{\mathcal{S}}$ production is less than 6.7% at 95% CL.

 $^{^5}$ Not seen in 19.7 fb $^{-1}$ of $p\,p$ collisions data at $\sqrt{s}=8$ TeV with $B_{\rm S}$ mesons reconstructed in decays to $J/\psi\,\phi$. An upper limit on the production rate times branching fraction for $X(5568)^\pm\to~B_{\rm S}\,\pi^\pm$ relative to inclusive $B_{\rm S}$ production is less than 1.1% at $p_T(B_{\rm S})>10$ GeV/c and less than 1.0% at $p_T(B_{\rm S})>15$ GeV/c at 95%CL.

⁶ Not seen in 3 fb⁻¹ of pp collision data at $\sqrt{s}=7$ and 8 TeV in a scan over the X(5568) mass and width, with B_S mesons reconstructed in decays to $D_S^-\pi^+$ or $J/\psi\phi$. An upper limit on the production rate times branching fraction for $X(5568)^\pm\to B_S\pi^\pm$ relative to inclusive B_S production is less than 2.1% at $p_T(B_S)>10$ GeV/c at 90% CL.